


YOUR ELIQUIS DOSING CALENDAR

For the treatment of deep vein thrombosis (DVT) and pulmonary embolism (PE) blood clots, and reducing the risk of them happening again after receiving treatment for blood clots in adults


If you're taking ELIQUIS **for the treatment of DVT/PE**, there is an important dosing change after 7 days. Use the calendar below to help keep track of your ELIQUIS dosage during the first 2 weeks of treatment. **Important note: treatment with ELIQUIS is likely to continue beyond 2 weeks.** Remember to always take ELIQUIS exactly as prescribed by your healthcare provider. Do not stop taking ELIQUIS without talking to the healthcare provider who prescribed it for you.

First 7 days



Day	1	2	3	4	5	6	7


After 7 days



Day	8	9	10	11	12	13	14

After 14 days, continue therapy as discussed with your healthcare provider.

After at least 6 months of DVT/PE treatment



If you're taking ELIQUIS to **reduce the risk of recurrence** after at least 6 months of initial treatment for DVT/PE, see the box on the left to learn about your extended therapy dose. Always be sure to take ELIQUIS exactly as prescribed by your healthcare provider.

Questions?

If you have questions, review your prescription with your healthcare provider to make sure you take the correct dosage strength and number of tablets.

(Please see the next page for how to take ELIQUIS.)

SELECTED IMPORTANT SAFETY INFORMATION

ELIQUIS can cause bleeding, which can be serious, and may lead to death.

Spinal or epidural blood clots (hematoma). People who take ELIQUIS, and have medicine injected into their spinal and epidural area, or have a spinal puncture have a risk of forming a blood clot that can cause long-term or permanent loss of the ability to move (paralysis).

Please see additional Important Safety Information on pages 2 and 3, and **U.S. Full Prescribing Information**, including **Boxed WARNINGS** and **Medication Guide**, or visit ELIQUIS.com.

How to Take ELIQUIS® (apixaban)



Be sure to take ELIQUIS exactly as prescribed by your healthcare provider. If you miss a dose of ELIQUIS, take it as soon as you remember on the same day, and do not take 2 doses at the same time to make up for a missed dose. Return to your usual dosing schedule the next day.



Do not take ELIQUIS if you currently have certain types of abnormal bleeding or have had a severe allergic reaction to ELIQUIS or any of the ingredients. Ask your healthcare provider if you're not sure.



Do not stop taking ELIQUIS without first talking with your healthcare provider. If you stop taking ELIQUIS, you may have increased risk of forming a clot in your blood.



Do not run out of ELIQUIS. Refill your prescription before you run out.

While taking ELIQUIS, you may bruise more easily and it may take longer than usual for any bleeding to stop.

If you take too much ELIQUIS, call your healthcare provider or go to the nearest hospital emergency room right away.

SELECTED IMPORTANT SAFETY INFORMATION

ELIQUIS may cause serious side effects, including:

- **Increased risk of blood clots if you stop taking ELIQUIS.** ELIQUIS lowers your chance of having a stroke by helping to prevent clots from forming. **Do not stop taking ELIQUIS without talking to the healthcare provider who prescribed it for you. Stopping ELIQUIS increases your risk of having a stroke.**

ELIQUIS may need to be stopped prior to surgery or a medical or dental procedure. Your healthcare provider will tell you when you should stop taking ELIQUIS and when you may start taking it again. If you have to stop taking ELIQUIS, your healthcare provider may prescribe another medicine to help prevent a blood clot from forming.

- **ELIQUIS can cause bleeding**, which can be serious, and may lead to death. This is because ELIQUIS is a blood thinner medicine that reduces blood clotting.

You may have a higher risk of bleeding if you take ELIQUIS and take other medicines that increase your risk of bleeding, such as aspirin, nonsteroidal anti-inflammatory drugs (called NSAIDs), warfarin, heparin, clopidogrel, selective serotonin reuptake inhibitors (SSRIs) or serotonin norepinephrine reuptake inhibitors (SNRIs), and other medicines to help prevent or treat blood clots.

Tell your healthcare provider about all the medicines you take, including any over-the-counter medicines, vitamins, and herbal supplements.

While taking ELIQUIS, you may bruise more easily and it may take longer than usual for any bleeding to stop.

Please see additional Important Safety Information on pages 1 and 3, and **U.S. Full Prescribing Information**, including **Boxed WARNINGS** and **Medication Guide**, or visit ELIQUIS.com.

Eliquis
(apixaban) tablets 5mg
2.5mg

SELECTED IMPORTANT SAFETY INFORMATION

ELIQUIS can cause bleeding (cont'd)

Call your healthcare provider or get medical help right away if you have any of these signs or symptoms of bleeding when taking ELIQUIS:

- unexpected bleeding or bruising, or bleeding that lasts a long time, such as unusual bleeding from the gums, nose bleeds that happen often, or menstrual or vaginal bleeding that is heavier than normal
 - bleeding that is severe or you cannot control
 - red, pink, or brown urine; red or black stools (looks like tar)
 - coughing up or vomiting blood or vomit that looks like “coffee grounds”
 - unexpected pain, swelling, or joint pain
 - headaches, or feeling dizzy or weak
- **Spinal or epidural blood clots (hematoma).** People who take ELIQUIS, and have medicine injected into their spinal and epidural area, or have a spinal puncture have a risk of forming a blood clot that can cause long-term or permanent loss of the ability to move (paralysis). Your risk of developing a spinal or epidural blood clot is higher if:
- a thin tube called an epidural catheter is placed in your back to give you certain medicine
 - you take NSAIDs or a medicine to prevent blood from clotting
 - you have a history of difficult or repeated epidural or spinal punctures
 - you have a history of problems with your spine or have had surgery on your spine

If you take ELIQUIS and receive spinal anesthesia or have a spinal puncture, your healthcare provider should watch you closely for symptoms of spinal or epidural blood clots or bleeding. Tell your healthcare provider or get medical help right away if you have back pain, tingling, numbness, or muscle weakness, especially in your legs and feet, or loss of control of the bowels or bladder (incontinence).

- ELIQUIS is not for use in people with artificial heart valves.
- ELIQUIS is not for use in people with antiphospholipid syndrome (APS), especially with positive triple antibody testing.

Do not take ELIQUIS if you currently have certain types of abnormal bleeding or have had a severe allergic reaction to ELIQUIS or any of the ingredients.

Before taking ELIQUIS, tell your healthcare provider about all your medical conditions, including if you have or ever had bleeding problems, have kidney or liver problems, or have antiphospholipid syndrome. Tell your healthcare provider right away if you are pregnant or breastfeeding, or plan to become pregnant or breastfeed. Taking ELIQUIS during pregnancy may increase the risk of bleeding in you or in your unborn baby. Do not breastfeed during treatment with ELIQUIS. **Females who are able to become pregnant:** talk with your healthcare provider about pregnancy planning, and your risk of severe uterine bleeding if you are treated with ELIQUIS.

Take ELIQUIS exactly as prescribed by your healthcare provider. Take ELIQUIS twice every day, and do not change your dose or stop taking it unless your healthcare provider tells you to. If you miss a dose of ELIQUIS, take it as soon as you remember on the same day, and do not take 2 doses at the same time to make up for a missed dose. **Do not stop taking ELIQUIS without first talking with your healthcare provider. Do not run out of ELIQUIS. Refill your prescription before you run out.** When leaving the hospital following hip or knee replacement, be sure that you will have ELIQUIS available to avoid missing any doses.

The most common side effect of ELIQUIS in adults was bleeding.

Please see additional Important Safety Information on pages 1 and 2, and **U.S. Full Prescribing Information**, including **Boxed WARNINGS** and **Medication Guide**, or visit [ELIQUIS.com](https://www.ELIQUIS.com).

Eliquis
(apixaban) tablets 5mg
2.5mg



ELIQUIS® and the ELIQUIS logo are trademarks of Bristol-Myers Squibb Company.
© 2025 Bristol-Myers Squibb Company. 432-US-2500134 05/25